

Which Scoring System is Better in Predicting Mortality in Multiple Trauma Patients: Revised Trauma Score or Glasgow Coma Scale

Adem Az, Çiğdem Orhan

Beylikdüzü State Hospital, Clinic of Emergency Medicine, İstanbul, Türkiye

Abstract

Objective: We investigated the prognostic value of the revised trauma score (RTS) and Glasgow Coma scale (GCS) in predicting mortality in multi-trauma patients.

Materials and Methods: This retrospective study included 537 consecutive trauma patients with a shock index ≥ 1.0 . We evaluated the demographics, clinical characteristics, and trauma scores, including GCS and RTS, in both the survivor and non-survivor groups.

Results: A total of 537 patients, comprising 58.29% males and 41.71% females, with a mean age of 44.46 ± 22.05 years, were included. Overall mortality was 13.04%. Age and sex differed significantly between survivors and non-survivors ($p=0.0001$ and $p=0.001$). Non-survivors had significantly lower mean GCS and RTS scores ($p=0.0001$ for both comparisons). Receiver operating characteristic analysis identified a $GCS \leq 10$ for predicting mortality in multi-trauma patients, with 99.89% sensitivity and 99.79% specificity. Additionally, an $RTS \leq 8$ had 98.57% sensitivity and 99.79% specificity for determining mortality.

Conclusion: Our results indicated that lower mean GCS and RTS scores were predictors of mortality in multi-trauma patients. A GCS of ≤ 10 and an RTS of ≤ 8 exhibited exceptional sensitivity and specificity for determining mortality in multi-trauma patients.

Keywords: Trauma, trauma scores, Glasgow Coma scale, revised trauma score, mortality

Introduction

Traumatic injuries represent a significant global health concern. Each year, more than 45 million people worldwide suffer from moderate to severe disabilities due to trauma. Furthermore, trauma-related injuries claim the lives of approximately 5.8 million individuals annually [1,2]. Moreover, 50%-60% of post-traumatic deaths occur within the initial hour [3]. Despite advances in healthcare and technology, fatalities in the scene or within the first hour persist as a significant public health issue. It is estimated that one-third of trauma-related deaths can be prevented with improved trauma systems [4].

In a study conducted in Türkiye, Höke et al. [5] investigated various trauma scores, including the injury severity score (ISS), new ISS, revised trauma score (RTS), and Glasgow Coma scale

(GCS), and observed that all of these scores demonstrated statistical significance in predicting mortality. In another study involving 633 trauma patients, Orhon et al. [6] found that GCS and RTS were significant indicators of mortality. Although numerous trauma scores are used to assess the severity of injuries and monitor clinical outcomes in trauma patients, the most accurate and reliable scoring system for determining morbidity and mortality remains unclear.

This study aimed to investigate the prognostic value of RTS and GCS in predicting mortality in patients with a shock index (SI) ≥ 1.0 who presented to the emergency department (ED) with multi-trauma.



Address for Correspondence: Adem Az, Beylikdüzü State Hospital, Clinic of Emergency Medicine, İstanbul, Türkiye

Phone: +90 530 100 71 17 E-mail: adem.aaz@gmail.com ORCID-ID: orcid.org/0000-0002-7204-6185

Received: 16.07.2023 Accepted: 14.09.2023



Copyright© 2024 The Author. Published by Galenos Publishing House on behalf of the Turkish Emergency Medicine Foundation. This is an open access article under the Creative Commons AttributionNonCommercial 4.0 International (CC BY-NC 4.0) License.

Materials and Methods

Ethics Committee Approval and Patient Consent

This study was conducted in accordance with the 1989 Declaration of Helsinki and was approved by the Institutional Review Board (IRB) of University of Health Sciences Türkiye, Haseki Training and Research Hospital Clinical Research Ethics Committee (approval number: 110-2022, date: 08.06.2022). As neither the images nor the accompanying text contained potentially identifying markers or patient identifiers, the IRB did not require patient consent for the review of their medical records.

Study Design and Setting

This retrospective, observational, single-center study included 537 consecutive adult patients (≥ 18 years old) who were admitted to our ED with multi-trauma between April 2021 and April 2022. Our hospital is one of the high-volume EDs in Istanbul, handling approximately 1500 emergency patient admissions daily. In addition, as a trauma center, our facility provides care for over 200 trauma patients daily, ranging from mild to severe cases. Only patients with a SI ≥ 1.0 were included in the study to exclude mild cases. The hospital's automation systems and archives were scanned for information on all patients presenting for the evaluation and treatment of acute traumatic injuries.

We assessed patients' demographics (age and sex), vital signs on admission [systolic blood pressure (SBP), respiratory rate, and heart rate (HR)], complaints and symptoms at admission, anatomic region of injury, type of trauma (blunt or penetrating), mechanism of injury, alcohol consumption, trauma scoring systems (GCS and RTS), and clinical outcomes (discharge, hospitalization, or death). In addition, SI was calculated for each patient. SI is defined as the ratio of HR to SBP.

Multi-trauma was defined as an injury to at least two body regions. Patients who experienced blunt or penetrating injuries in the same anatomical region were classified as having penetrating injuries. This study classified multiple injuries to the same anatomical region as a singular injury to that specific anatomical region.

The patients in the study cohort were categorized into survivors and non-survivors. Demographics, clinical characteristics, and trauma scores (GCS and RTS) were compared among the groups to determine the factors associated with mortality.

Study Population and Sampling

All cases meeting the eligibility criteria were included to prevent selection bias. We enrolled 6,978 patients admitted to the ED due to traumatic injuries between April 2021 and April 2022. Patients with non-traumatic injuries or those presenting to the ED for any other reason were subsequently excluded. Additionally, 152 patients were excluded because

of a lack of information. Moreover, 2,348 patients under the age of 18 years were excluded from the study. Furthermore, 1,926 patients were excluded because they had mono-trauma. Moreover, 2015 patients with a SI < 1.0 were excluded because of severe injuries. The remaining 537 patients were included in the study (Figure 1).

Trauma Assessment Scores

GCS is a neurological assessment tool that measures a person's level of consciousness based on eye-opening, verbal, and motor responses, which are assigned 4, 5, and 6 points, respectively (for a total score of 15 points).

The RTS is a tool used to assess the severity of a traumatic injury. It considers three key parameters: GCS, SBP, and respiratory rate, with a total score of 12 points.

Statistical Analysis

All data analyses were conducted using SPSS statistical software (version 15.0 for Windows; SPSS Inc., Chicago, IL, USA). Categorical variables are expressed as numbers of patients (n) and percentages (%). Numerical data are expressed as mean, standard deviation, and minimum and maximum values. Intragroup analyses (survivors vs non-survivors) were conducted using the chi-square test for normally distributed data and the Mann-Whitney U-test for non-normally distributed data. Independent variables predicting mortality (age, sex, GCS and RTS) were analyzed using multivariate logistic regression analysis. The receiver operating characteristic (ROC) curve was used to determine the cut-off point for GCS and RTS. The threshold for statistical significance was defined as $p < 0.05$.

Results

The demographic and clinical characteristics of the trauma patients are presented in Table 1. The study comprised a sample size of 537 patients, with 313 (58.29%) males and 224 (41.71%) females. The mean age was 44.46 ± 22.05 years, with a range of 18-96 years. The overall mortality rate was 13.04%. In addition, 26.82% of the patients were discharged from the ED, and 61.64% were hospitalized. Overall, 73.93%




Figure 1. Flowchart

of patients presented with blunt injuries. Falls were the most commonly reported mechanism of trauma, accounting for 52.51%, followed by traffic accidents at 17.13%, and accidental injuries at 11.17%. A total of 64 individuals were transported to the ED via ambulance, while 473 arrived on foot. Analysis of anatomical regions affected by injuries revealed that the head and face were the most prevalent sites, comprising 43.58% of the cases.

A total of 142 patients were hospitalized and followed up in the orthopedics department. Additionally, 64 individuals

received treatment in the neurosurgery department, 54 in the general surgery department, 30 in the cardiovascular surgery department, 22 in the thoracic surgery department, and 8 in the intensive care unit.

Table 2 presents the comparative analysis of demographics, clinical characteristics, and trauma scores among patients who survived and those who did not. The age of non-survivors was found to be significantly lower than that of survivors ($p=0.0001$). Furthermore, the prevalence of males was significantly higher among non-survivors than among survivors ($p=0.001$). Penetrating traumas occurred significantly more commonly in non-survivors than in survivors ($p=0.024$). Moreover, statistically significant differences were observed among non-survivors and survivors in terms of the mechanisms of trauma such as falls, traffic accidents, assault, and gunshot wounds ($p=0.0001$, $p=0.014$, $p=0.002$, and $p=0.001$, respectively). Finally, non-survivor patients had significantly lower mean GCS and RTS scores than survivors ($p=0.0001$ for both comparisons).

Multivariate logistic regression analysis demonstrated that increased age [odds ratio (OR): 0.98, 95% confidence interval

Table 1. Demographic and clinical characteristics of trauma patients

Characteristics			
Age in years, mean \pm SD (min - max)		44.46 \pm 22.05 (18-96)	
		n	(%)
Sex	Female	224	(41.71)
	Male	313	(58.29)
The types of trauma	Blunt	397	(73.93)
	Penetrating	140	(26.07)
Mechanism of trauma	Fall	282	(52.51)
	Traffic accident	92	(17.13)
	Assault	60	(11.17)
	Accidental injuries	51	(9.50)
	Stab wounds	34	(6.33)
	Gunshot wounds	18	(3.35)
Place of trauma	Street/road/highway	351	(65.36)
	Home	103	(19.18)
	Commercial/work	83	(15.46)
Alcohol consumed	No	420	(78.21)
	Yes	117	(21.79)
Forensic trauma	No	186	(34.64)
	Yes	351	(65.36)
Transport to the hospital	By foot	473	(88.08)
	Via ambulance	64	(11.92)
Anatomic region of injury	Head and face	234	(43.58)
	Lower extremities	222	(41.34)
	Upper extremities	201	(37.43)
	Abdomen	191	(35.57)
	Chest	184	(34.26)
	Spine	151	(28.12)
Outcome	Discharge	144	(26.82)
	Hospitalization	331	(61.64)
	Death	70	(13.04)

Data are given as numbers (n) and percentages (%), mean, standard deviation (SD), and minimum and maximum values
min - max: Minimum - maximum

Table 2. Comparison of demographic characteristics and trauma scores between patients who survived and those who did not

	Survivors		Non-survivors		p
Age in years, mean \pm SD	46.05 \pm 22.34		33.83 \pm 16.54		0.0001
	n	(%)	n	(%)	p
Sex					
Female	208	(44.54)	16	(22.86)	0.001
Male	259	(55.46)	54	(77.14)	
Mechanism of trauma					
Fall	264	(56.53)	19	(27.14)	0.0001
Traffic accident	73	(15.63)	18	(25.71)	0.014
Accidental injuries	47	(10.06)	4	(5.71)	0.348
Assault	44	(9.42)	16	(22.86)	0.002
Stab wounds	29	(6.21)	5	(7.14)	0.972
Gunshot wounds	10	(2.14)	8	(11.43)	0.001
The types of trauma					
Blunt	353	(75.59)	44	(62.86)	0.024
Penetrating	114	(24.41)	26	(37.14)	
	Mean \pm SD		Mean \pm SD		p
Glasgow Coma scale	14.95 \pm 0.24		5.04 \pm 2.07		0.0001
Revised trauma score	11.94 \pm 0.27		5.61 \pm 1.83		0.0001

Data are given as numbers (n) and percentages (%), mean, and standard deviation (SD)
*Intragroup analyses (survivors vs non-survivors) were conducted using the chi-square test for normally distributed data and the Mann-Whitney U-test for non-normally distributed data, as appropriate

(CI): 0.96-1.01; $p=0.001$], female gender (OR: 1.96, 95% CI: 1.06-2.61; $p=0.031$), and decreased GCS (OR: 0.64, 95% CI: 0.18-0.98; $p=0.027$) and RTS scores (OR: 0.64, 95% CI: 0.22-1.97; $p=0.049$) were identified as significant predictors of mortality among trauma patients (Table 3).

ROC analysis identified a GCS cut-off score of ≤ 10 to determine mortality in multi-trauma patients, with 99.89% sensitivity and 99.79% specificity [area under the curve (AUC): 0.999, 95% CI 0.991-0.999; Table 4 and Figure 2]. In addition, ROC analysis revealed a cut-off RTS of ≤ 8 , with 98.57% sensitivity and 99.79% specificity for determining mortality in multi-trauma patients (AUC: 0.99, 95% CI 0.990-1.000; Table 4 and Figure 2).

Discussion

Trauma is one of the leading causes of mortality. Annually, trauma leads to the mortality of nearly 6 million individuals worldwide [1]. A substantial number of fatalities occur either at the scene of the incident or within the initial 4 h following the patient’s arrival at an ED [2]. Hence, the main goal of this study was to predict and determine individuals at an increased risk of mortality at an early stage. The key findings of our study are as follows. First, males and young adults exhibited a higher prevalence of trauma and trauma-related mortality. Second, falls, traffic accidents, and accidental injuries were the most commonly reported mechanisms of trauma. Third, non-survivors had lower mean GCS and RTS scores than survivors. Fourth, in determining mortality in multi-trauma patients, a GCS score of ≤ 10 was found to be the cut-off with 99.89% sensitivity and 99.79% specificity, and an RTS score of ≤ 8 was determined as the cutoff with 98.57% sensitivity and 99.79% specificity.

In studies analyzing the epidemiologic and demographic features of trauma patients, Mutasingwa and Aaro [7] and

Aluisio et al. [8] consistently noted that young males were more commonly presented to the ED with traumatic injuries. Additionally, in the United States, trauma is the leading cause of mortality among individuals under the age of 44 [9]. Similarly, in our study, males and young adults exhibited a higher prevalence of trauma and trauma-related mortality.

According to our findings, the prevailing causes of trauma were falls and traffic accidents. Consistent with our study, Chokotho et al. [10] reported that falls and traffic accidents were the most common mechanisms of injury in their study involving 49,241 trauma cases. In another study conducted in Türkiye, Çırak et al. [11] found that falls and traffic accidents were the leading causes of trauma among patients. In studies conducted in low- or middle-income countries, Rouhani et al., [12], Soundarrajan et al., [13], and Zuraik and Sampalis [14] discovered that road traffic accidents were the most common trauma mechanism, followed by falls. Our findings are consistent with the main causes of trauma worldwide. However, the prevalence and trends of trauma may vary across various cultural contexts, nations, and socioeconomic circumstances. The higher incidence of traffic accidents, particularly in low- and middle-income countries, can be




Figure 2. Specificity and sensitivity of GCS and RTS scores for determining mortality in multi-trauma patients using receiver operating characteristic curves [area under the curve (AUC): 0.999, 95% confidence interval (CI) 0.991-0.999 and AUC: 0.99, 95% CI 0.990-1.000; respectively]

GCS: Glasgow Coma scale, RTS: Revised trauma score

Table 3. Multivariate logistic regression analysis to determine mortality				
	p	OR	95% CI	
Age in years	0.001	0.98	0.96	1.01
Sex (female)	0.031	1.96	1.06	2.61
Glasgow Coma scale	0.027	0.64	0.18	0.98
Revised trauma scores	0.049	0.64	0.22	1.97

OR: Odds ratio, CI: Confidence interval

Table 4. Trauma scores for determining mortality in multi-trauma patients								
Criterion	AUC	SE	95% CI	Sensitivity	Specificity	PPV	NPV	LR (+)
GCS ≤ 10	0.999	0.001	0.991-0.999	99.89	99.79	98.6	100.0	467.00
RTS ≤ 8	0.999	0.001	0.990-1.000	98.57	99.79	98.7	99.8	460.33

AUC: Area under the curve, SE: Standard error, CI: Confidence interval, PPV: Positive predictive value, NPV: Negative predictive value, LR (+): Likelihood ratio, GCS: Glasgow Coma scale, RTS: Revised trauma score

attributed to inadequate adherence to safety precautions and less compliance with traffic regulations [10,14]. The research conducted within Türkiye revealed a higher prevalence of fall incidents compared with traffic accidents [11]. Moreover, based on our findings, penetrating injuries exhibited a higher fatality rate, even though most trauma incidents involved blunt injuries.

In a study involving a sample of 633 trauma patients from Türkiye, Orhon et al. [6] found that GCS and RTS were significant indicators of mortality. In another study conducted in Türkiye, Güneytepe et al. [15] investigated various trauma scores, including GCS, RTS, ISS, and trauma and injury severity score (TRISS), and observed that all these scores demonstrated statistical significance in predicting mortality. In a study of 1,410 trauma patients, Yadollahi et al. [16] also observed that TRISS, RTS, GCS, and ISS were all highly effective in determining prognosis and mortality among trauma patients. Similarly, our study revealed significant differences between survivors and non-survivors in terms of GCS and RTS.

In a study assessing post-traumatic deaths, Demetriades et al. [3] discovered that GCS <8 emerged as the most important risk factor associated with mortality among trauma patients within the first hour after admission to the hospital. Another study, involving 740 trauma patients, also recognized a GCS <8 as a reliable predictor of mortality [17]. Furthermore, Yadollahi et al. [16] identified increased age, GCS <8 , RTS <7.6 , and TRISS <0.9 as the most significant predictors of in-hospital mortality. Our results demonstrated that GCS could predict mortality with 99.89% sensitivity and 99.79% specificity in the scores ≤ 10 . Moreover, patients with a GCS of ≤ 10 have a 467-fold increased risk of mortality than those with a GCS of >10 . Similar to our findings, a study conducted in Northern Iran reported that a GCS ≤ 8 predicts mortality with exceptionally high accuracy, showing a sensitivity of 98.4% and specificity of 92.3% [18].

Yadollahi et al. [16] demonstrated that RTS exhibited the highest effectiveness in assessing the severity of traumatic injuries, following TRISS. Furthermore, they established a cut-off point for RTS at ≤ 7.69 with 95% sensitivity and 67% specificity in predicting mortality in trauma patients. In another study conducted by Yousefzadeh-Chabok et al. [19], an RTS score of ≤ 6 was identified as a predictor of mortality among trauma patients, exhibiting 99% sensitivity and 62% specificity. In our study, an RTS score of ≤ 8 was determined as a predictor of mortality with 98.57% sensitivity and 99.79% specificity. In our cohort, we exclusively included patients with SI ≥ 1.0 . The higher specificity observed in our findings compared with other studies can be attributed to this selection criterion. Based on our findings, the combined use of RTS with SI offers valuable insights for predicting mortality and prognosis among multi-trauma patients.

In our multivariate logistic regression analysis that examined the utility of the GCS and RTS for predicting mortality in multi-trauma patients, we found that both scoring systems had a comparable OR. However, the GCS showed a slightly higher level of statistical significance and a more reliable CI, suggesting that it may be a more reliable predictor of mortality in our population.

Study Limitations

A limitation of this study is its use of a retrospective and hospital-based study design, which offers a risk of selection and misclassification biases affecting the obtained results. Second, our observations are limited to the patient population that seeks medical attention at the hospital. Consequently, it is not possible to reach conclusions about the prevalence of trauma among the general population. Finally, we lack information about the post-discharge health status and care quality of trauma patients.

Conclusion

Our results indicated that lower mean GCS and RTS scores were predictors of mortality in multi-trauma patients. Specifically, a GCS of ≤ 10 had a sensitivity of 99.89% and a specificity of 99.79% for determining mortality in multi-trauma patients with an SI ≥ 1.0 . Moreover, an RTS of ≤ 8 exhibited an exceptional sensitivity of 98.57% and a specificity of 99.79% in identifying mortality. We recommend the use of trauma scores, such as GCS and RTS, in conjunction with SI at ED admission to accurately assess disease severity and mortality risk in trauma patients.

Ethics

Ethics Committee Approval: This study was conducted in accordance with the 1989 Declaration of Helsinki and was approved by the Institutional Review Board (IRB) of University of Health Sciences Türkiye, Haseki Training and Research Hospital Clinical Research Ethics Committee (approval number: 110-2022, date: 08.06.2022).

Informed Consent: As neither the images nor the accompanying text contained potentially identifying markers or patient identifiers, the IRB did not require patient consent for the review of their medical records.

Authorship Contributions

Concept: A.A., Ç.O., Design: A.A., Ç.O., Data Collection or Processing: A.A., Ç.O., Analysis or Interpretation: A.A., Literature Search: A.A., Ç.O., Writing: A.A., Ç.O.

Conflict of Interest: No conflicts of interest were declared by the authors.

Financial Disclosure: The authors declare that this study received no financial support.

References

1. World Health Organization. The global burden of disease: 2004 update 2008. Geneva; 2008. Available from: <https://apps.who.int/iris/handle/10665/43942>
2. World Health Organization. Injuries and violence: the facts. Geneva; 2010. Available from: <https://www.who.int/news-room/fact-sheets/detail/injuries-and-violence>
3. Demetriades D, Murray J, Charalambides K, Alo K, Velmahos G, Rhee P, et al. Trauma fatalities: time and location of hospital deaths. *J Am Coll Surg.* 2004;198:20-6.
4. Mock C, Joshipura M, Arreola-Risa C, Quansah R. An estimate of the number of lives that could be saved through improvements in trauma care globally. *World J Surg.* 2012;36:959-63.
5. Höke MH, Usul E, Özkan S. Comparison of Trauma Severity Scores (ISS, NISS, RTS, BIG Score, and TRISS) in Multiple Trauma Patients. *J Trauma Nurs.* 2021;28:100-6.
6. Orhon R, Eren SH, Karadayı S, Korkmaz I, Coşkun A, Eren M, et al. Comparison of trauma scores for predicting mortality and morbidity on trauma patients. *Ulus Travma Acil Cerrahi Derg.* 2014;20:258-64.
7. Mutasingwa DR, Aarø LE. Injury registration in a developing country. A study based on patients' records from four hospitals in Dar es Salaam, Tanzania. *Cent Afr J Med.* 2001;47:203-9.
8. Aluisio AR, De Wulf A, Louis A, Bloem C. Epidemiology of Traumatic Injuries in the Northeast Region of Haiti: A Cross-sectional Study. *Prehosp Disaster Med.* 2015;30:599-605.
9. Centers for Disease Control and Prevention. Injuries and Violence Are Leading Causes of Death. Available from: <https://www.cdc.gov/injury/wisqars/animated-leading-causes.html>
10. Chokotho L, Croke K, Mohammed M, Mulwafu W, Bertfelt J, Karpe S, et al. Epidemiology of adult trauma injuries in Malawi: results from a multisite trauma registry. *Inj Epidemiol.* 2022;9:14.
11. Çırak B, Güven MB, Işık S, Kıymaz N, Demir Ö. An epidemiologic study an Patients admitted to emergency service. *Ulus Travma Acil Cerrahi Derg.* 1999;5:157-9.
12. Rouhani SA, Eliacin HC, Edmond MC, Checkett KA, Rimpel L, Marsh RH. Epidemiology of traumatic injuries presenting to an ED in Central Haiti: a retrospective cohort study. *Emerg Med J.* 2019;36:389-94.
13. Soundarrajan A, Jena NN, Smith J, Douglass K. 165 A Retrospective Observational Study on Epidemiology of Traumatic Injuries Presenting to a Tertiary Care Hospital in Madurai, India. 2017;70:s66. [chrome-extension://efaidnbmnnnibpcajpcglclefindmkaj/https://www.annemergmed.com/action/showPdf?pii=S0196-0644%2817%2931092-2](https://www.annemergmed.com/action/showPdf?pii=S0196-0644%2817%2931092-2)
14. Zuraik C, Sampalis J. Epidemiology of Traumatic Injuries at an Urban Hospital in Port-au-Prince, Haiti. *World J Surg.* 2017;41:2674-80.
15. Güneytepe Üİ, Aydın ŞA, Gökçöz Ş, Özgüç H, Ocakoğlu G, Aktaş H. The factors influencing the mortality in elderly trauma patients and scoring systems. *Uludag Medical Journal.* 2008;34:15-9.
16. Yadollahi M, Kashkooe A, Rezaiee R, Jamali K, Niakan MH. A Comparative Study of Injury Severity Scales as Predictors of Mortality in Trauma Patients: Which Scale Is the Best? *Bull Emerg Trauma.* 2020;8:27-33.
17. Jelodar S, Jafari P, Yadollahi M, Sabetian Jahromi G, Khalili H, Abbasi H, et al. Potential Risk Factors of Death in Multiple Trauma Patients. *Emerg (Tehran).* 2014;2:170-3.
18. Norouzi V, Feizi I, Vatankhah S, Pourshaikhian M. Calculation of the probability of survival for trauma patients based on trauma score and the injury severity score model in fatemi hospital in ardabil. *Arch Trauma Res.* 2013;2:30-5.
19. Yousefzadeh-Chabok S, Hosseinpour M, Kouchakinejad-Eramsadati L, Ranjbar F, Malekpouri R, Razzaghi A, et al. Comparison of Revised Trauma Score, Injury Severity Score and Trauma and Injury Severity Score for mortality prediction in elderly trauma patients. *Ulus Travma Acil Cerrahi Derg.* 2016;22:536-40.